ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 69 • Numbers 1–4 • 2002

Author index

ABU ELZEIN, E.M.E.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

ADEJINMI, J.O.

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

AL NAIEM, A.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

AL-AFALEQ, A.I.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

AL-ATIYA, S.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

ALBERTYN, J.

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

ALLSOPP, B.A.

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

ALLSOPP, MARIA T.E.P.

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

ALMLI, B.

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

ATANÁSIO, A.

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

BENGIS, R.G.

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

BETT, B.

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

BOOMKER, J.

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

BOORMAN, J.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

BOSMAN, ANNA-MARI

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

BOTHA, C.J.

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

BRAGG, R.R.

Isolation of serovar C-3 *Haemophilus paragallinarum* from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of *H. paragallinarum*—R.R. BRAGG 129–132

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 1: NAD-dependent field isolates—R.R. BRAGG 163–169

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG 171–175

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

BRETT, O.L.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

BRYSON, N.R.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

BUBB, M.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

BURGESS, G.W.

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS 237–242

CHAMUNORWA, J.P.

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

CLAASEN, B.

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

COETZER, J.A.W.

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

DE KLERK, LIN-MARI

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

DIPEOLU, O.O.

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

DUNGU, B.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

EMSLIE, F.R.

An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL 123–127

ERASMUS, B.J.

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

ESPACH, A.

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

FLÅØYEN, A.

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

FOURIE, L.J.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

GEHRING, R.

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

GROBLER, D.G.

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

GROENEWALD, DALEEN

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

GUTHRIE, A.J.

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

HADGU, K.

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

HARRIS, H.J.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

HERING-HAGENBECK, S.F.B.N.

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

HILALI, M.A.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

HLATSHWAYO, M.

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

HOLTUG, K.

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

HORAK, I.G.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

HOWELL, P.G.

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

HUNTER, P.

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

JOSEMANS, ANTOINETTE I.

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

JUNKER, K.

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

KAPAGA, A.M.

Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA 305–314

KARANJA, S.M.

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

KELLERMAN, T.S.

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

KIDANEMARIAM, A.

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

KITHUKA, J.M.

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

KIVARIA, F.M.

Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA 305–314

KOK, D.J.

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

KRECEK, R.C.

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

KUMBA, F.F.

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

LABUSCHAGNE, L.

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

LOPEZ-REBOLLAR, LAURA M.

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

MAGONA, J.W.

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

MAINGI, N.

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

MAYENDE, J.S.P.

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

MBATI, P.A.

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

MEINTJES, T.

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

MEKONNEN, S.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

MELLOR, P.S.

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

MEYER, P.

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

MICHEL, ANITA L.

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

MUMBI, V.M.

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

MUSARA, C.

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

MUSONDA, M.M.

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

MWASE, M.

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

NEL, J.R.

An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL 123–127

NEL, L.H.

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

NGAIRA, J.M.

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

NGOTHO, J.W.

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

NJERUH, F.M.

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

OLUWANIYI, J.O.

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

OMBUI, J.N.

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

OTIENO, R.O.

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

PAWESKA, J.T.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

PETER, R.J.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

PETRUS, P.

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

PETTER, A.J.

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

POTTS, A.D.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

PROZESKY, L.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

ROMITO, M.

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

SAHLE, M.

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS 237–242

SCHULTZ, R.A.

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

SITOE, C.

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

SIVERTSEN, T.

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

SKADHAUGE, E.

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

SMITH, S.J.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

SNYMAN, L.D.

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

SPICKETT, A.M.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

STRYDOM, T.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

TAIWO, V.O.

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

TAOLE, M.

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

TAYLOR, R.J.

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

VAN HEERDEN, E.

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

VAN HEERDEN, HENRIETTE

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

VAN ROOYEN, J.M.

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

VAN STRIJP, M. FRANSI

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

VAN WYK, LAETITIA

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

VAN WYK, J.A.

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

VAN ZYL, W.

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

VATTA, A.F.

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

VENTER, D.

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

VILJOEN, G.J.

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

VISAGE, CARINA W.

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

VORSTER, J.H.

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

WARUIRU, R.M.

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

WEDA, E.H.

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

ZWEYGARTH, E.

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA, M. LOPEZ-REBOLLAR and P. MEYER 197–200

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 69 • Numbers 1–4 • 2002

Subject index

Acari: Ixodidae

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

Acaricide

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

African buffaloes

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

Albendazole

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

Amphistome infection

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

Anaplasmosis

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

Animal health

Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA 305–314

Anthelmintic resistance

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Antigen preparation

Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS 237–242

Arbovirus

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi

Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

Ateleriz frontalis

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

Babesia caballi

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Boophilus decoloratus

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

Bovine fasciolosis

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

Bovine tuberculosis

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

Brucella abortus

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

Brucella melitensis

An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL 123–127

Bufadienolide

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Camels

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

Cape mountain zebra

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Cardiac glycosides

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Cattle

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

Chelonia

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

Closantel

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

Colon

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

Complement fixation test

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

Coryza

Isolation of serovar C-3 *Haemophilus paragallinarum* from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of *H. paragallinarum*—R.R. BRAGG 129–132

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

Cotyledoside

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Cryopreservation

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

Culicoides

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

Cycloheximide

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

Diptera: Ceratopogonidae

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

Dourine

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

Ehrlichia (Cowdria) ruminantium

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

Elephantulus myurus

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

Epoxyscillirosidin

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Equine encephalosis virus

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Equine piroplasmosis

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Equus zebra zebra

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Ethiopia

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

Faecal egg counts

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

Fasciola

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

Feed aversion

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

Fenbendazole

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANASIO, J. BOOMKER and C. SITOE 215–220

Gamma-interferon test

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

***Geigeria ornativa* (vermeerbos)**

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

Gekkonidae

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Gerrhosauridae

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Goats

An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL 123–127

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

Haematology

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

Haemonchus contortus

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

Haemophilus paragallinarum

Isolation of serovar C-3 *Haemophilus paragallinarum* from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of *H. paragallinarum*—R.R. BRAGG 129–132

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 1: NAD-dependent field isolates—R.R. BRAGG 163–169

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG 171–175

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

Heartwater

The Kümm isolate of *Ehrlichia ruminantium: In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

Helminthosis

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

Hippopotami

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

Horses

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

Indirect ELISA

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS 237–242

Information system

Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA 305–314

Kenya

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

Kruger National Park

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

Kümm isolate

The Kümm isolate of *Ehrlichia ruminantium*: *In vitro* isolation, propagation and characterization—E. ZWEYGARTH, ANTOINETTE I. JOSEMANS, M. FRANSI VAN STRIJP, HENRIETTE VAN HEERDEN, MARIA T.E.P. ALLSOPP and B.A. ALLSOPP 147–153

KwaZulu-Natal

An overview of the eradication of *Brucella melitensis* from KwaZulu-Natal—F.R. EMSLIE and J.R. NEL 123–127

Levamisole

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Lizards

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Madin Derby bovine kidney cells (MDBK)

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

McMaster technique

Freezing of sheep faeces invalidates *Haemonchus contortus* faecal egg counts by the McMaster technique—J.A. VAN WYK and LAETITIA VAN WYK 299–304

Methacholine

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Mites

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

Mozambique

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Mycobacterium bovis

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

NAD-dependence/NAD-independence

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 1: NAD-dependent field isolates—R.R. BRAGG 163–169

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG 171–175

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 3: experimentally produced NAD-independent isolate—M. TAOLE, J. ALBERTYN, E. VAN HEERDEN and R.R. BRAGG 189–196

Namibia

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

Nematodes

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Seasonal availability of gastrointestinal nematode larvae to cattle on pasture in the central highlands of Kenya—R.M. WARUIRU, E.H. WEDA, R.O. OTIENO and J.W. NGOTHO 141–146

A survey on the occurrence of resistance to anthelmintics of gastrointestinal nematodes of goats in Mozambique—A. ATANÁSIO, J. BOOMKER and C. SITOE 215–220

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

Neutralizing antibody

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Neutrophils

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

Newcastle disease virus

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

Development of solid phase antigen for indirect ELISA for the detection of specific antibody responses to infection with Newcastle disease virus—M. SAHLE and G.W. BURGESS 237–242

Nigeria

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

Orbivirus

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Osmosis

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

Ostrich

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

Oxyuroidea

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Pachydactylus

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground,

1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Parapharyngodon

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Pelomedusa

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

Pelonia africana

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

Pelusios

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

Pentastomida

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

Pharyngodonidae

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Plant poisoning

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

Psoroptes ovis

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

Rat jejunum

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Rhipicentor nuttalli

Hosts, seasonal occurrence and life cycle of *Rhipicentor nuttalli* (Acari: Ixodidae)—L.J. FOURIE, I.G. HORAK, D.J. KOK and W. VAN ZYL 177–187

***Rhipicephalus* spp.**

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

Rift Valley fever virus

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

RT-PCR

Development of a diagnostic one-tube RT-PCR for the detection of Rift Valley fever virus—A. ESPACH, M. ROMITO, L.H. NEL and G.J. VILJOEN 247–252

Saudi Arabia

Seasonal abundance of four *Culicoides* spp. (Diptera: Ceratopogonidae) at Al-Ahsa oasis, Eastern Province, Saudi Arabia—E.M.E. ABU ELZEIN, M.A. HILALI, A.I. AL-AFALEQ, P.S. MELLOR, J. BOORMAN, S. AL-ATIYA and A. AL NAIEM 115–122

Scincidae

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Sera

Validation of an indirect enzyme-linked immunosorbent assay for the detection of antibody against *Brucella abortus* in cattle sera using an automated ELISA workstation—J.T. PAWESKA, A.D. POTTS, H.J. HARRIS, S.J. SMITH, G.J. VILJOEN, B. DUNGU, O.L. BRETT, M. BUBB and L. PROZESKY 61–77

Serotypes

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Sheep

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

Strategic control of gastrointestinal nematodes of sheep in the highlands of central Kenya—N. MAINGI, R.O. OTIENO, V.M. MUMBI and E.H. WEDA 229–235

On-host ecology and off-host survival of the sheep scab mite *Psoroptes ovis*—T. MEINTJES, L.J. FOURIE and I.G. HORAK 273–283

Effects of the scab mite *Psoroptes ovis* on the haematology and live mass of Merino and Dorper sheep—T. MEINTJES, L.J. FOURIE and I.G. HORAK 285–293

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

Skrjabinodon

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

South Africa

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Description of *Pelonia africana* n. g., n. sp. (Pentastomida: Sebekidae) from the lungs of *Pelomedusa subrufa* and *Pelusios sinuatus* (Chelonia) in South Africa—K. JUNKER and J. BOOMKER 53–59

The classification of seven serotypes of equine encephalosis virus and the prevalence of homologous antibody in horses in South Africa—P.G. HOWELL, DALEEN GROENEWALD, CARINA W. VISAGE, ANNA-MARI BOSMAN, J.A.W. COETZER and A.J. GUTHRIE 79–93

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 1: NAD-dependent field isolates—R.R. BRAGG 163–169

Virulence of South African isolates of *Haemophilus paragallinarum*. Part 2: Naturally occurring NAD-independent field isolates—R.R. BRAGG 171–175

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

Spauligodon

Redescription of some *Spauligodon* spp. and *Parapharyngodon* spp., and of *Skrjabinodon mabuyae* Sandground, 1936) Inglis, 1968 (Pharyngodonidae: Oxyuroidea) from insectivorous South African lizards—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 7–29

Struthio camelus

Water absorption in relation to fermentation in the colon of the ostrich (*Struthio camelus*)—C. MUSARA, J.P. CHAMUNORWA, K. HOLTUG and E. SKADHAUGE 315–320

Syncerus caffer

The gamma-interferon test: its usefulness in a bovine tuberculosis survey in African buffaloes (*Syncerus caffer*) in the Kruger National Park—D.G. GROBLER, ANITA L. MICHEL, LIN-MARI DE KLERK and R.G. BENGIS 221–227

Tachygonetria

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Tanzania

Review of current problems and shortcomings in the Tanzanian animal health information system with suggestions on improvement—F.M. KIVARIA and A.M. KAPAGA 305–314

Teratogenicity

Teratogenicity of a mutagenised Rift Valley fever virus (MVP 12) in sheep—P. HUNTER, B.J. ERASMUS and J.H. VORSTER 95–98

Theileria equi

In vitro isolation of equine piroplasms derived from Cape mountain zebra (*Equus zebra zebra*) in South Africa—E. ZWEYGARTH, LAURA M. LOPEZ-REBOLLAR and P. MEYER 197–200

Theileriosis

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

Thelandros

Redescription of some *Thelandros* and *Tachygonetria* spp. (Pharyngodonidae: Oxyuroidea) from the omnivorous plated lizard, *Gerrhosaurus validus validus* A. Smith, 1849 in South Africa—S.F.B.N HERING-HAGENBECK, A.J. PETTER and J. BOOMKER 31–51

Ticks

Seasonal abundance of adult ixodid ticks infesting cattle belonging to resource-limited farmers in the north-eastern Free State Province of South Africa—M. HLATSHWAYO, P.A. MBATI and O.O. DIPEOLU 1–6

Acaricide resistance profiles of single- and multi-host ticks from communal and commercial farming areas in the Eastern Cape and North-West Provinces of South Africa—S. MEKONNEN, N.R. BRYSON, L.J. FOURIE, R.J. PETER, A.M. SPICKETT, R.J. TAYLOR, T. STRYDOM and I.G. HORAK 99–105

Trematode

The prevalence and economic importance of bovine fasciolosis in Kenya—an analysis of abattoir data—J.M. KITHUKA, N. MAINGI, F.M. NJERUH and J.N. OMBUI 255–262

Amphistome infection of goats farmed under resource-poor conditions in South Africa—A.F. VATTA and R.C. KRECEK 327–329

***Trypanosoma*/trypanosomosis**

Parasitological prevalence of bovine trypanosomosis in Kindo Koisha district, Wollaita zone, south Ethiopia—A. KIDANEMARIAM, K. HADGU and M. SAHLE 107–113

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

Non-immune control of trypanosomosis: *In vitro* oxidative burst of PMA- and trypanosome-stimulated neutrophils of Boran and N'Dama cattle—V.O. TAIWO, J.O. ADEJINMI and J.O. OLUWANIYI 155–161

Animal-level risk factors for *Trypanosoma evansi* infection in camels in eastern and central parts of Kenya—J.M. NGAIRA, B. BETT and S.M. KARANJA 263–271

Apparent prevalence of dourine in the Khomas region of Namibia—F.F. KUMBA, B. CLAASEN and P. PETRUS . 295–298

Tyledoside D

The effect of three bufadienolide cardiac glycosides on contraction of isolated rat jejunum—C.J. BOTHA, R. GEHRING, J.M. VAN ROOYEN and D. VENTER 243–246

Uganda

Occurrence of concurrent trypanosomosis, theileriosis, anaplasmosis and helminthosis in Friesian, Zebu and Sahiwal cattle in Uganda—J.W. MAGONA and J.S.P. MAYENDE 133–140

V4 strain replication

Multiplication of the V4 strain of Newcastle disease virus in Madin Derby bovine kidney cells—M. SAHLE, G.W. BURGESS and A. KIDANEMARIAM 201–206

Vaccines

Isolation of serovar C-3 *Haemophilus paragallinarum* from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of *H. paragallinarum*—R.R. BRAGG 129–132

Vermeersiekte

Continuous exposure to an aversive mixture as a means of maintaining aversion to vermeerbos (*Geigeria ornativa* O. Hoffm.) in the presence of non-averted sheep—L.D. SNYMAN, R.A. SCHULTZ, T.S. KELLERMAN and L. LABUSCHAGNE 321–325

Zambia

Hepatic and renal concentrations of copper and other trace elements in hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Kafue and Luangwa Rivers in Zambia—M. MWASE, B. ALMLI, T. SIVERTSEN, M.M. MUSONDA and A. FLÅØYEN 207–214

Zimbabwe

Isolation of serovar C-3 *Haemophilus paragallinarum* from Zimbabwe: A further indication of the need for the production of vaccines against infectious coryza containing local isolates of *H. paragallinarum*—R.R. BRAGG 129–132